



RRB

Junior Engineers (JE)

Railway Recruitment Board

Volume - 1

Mathematics




Table of Content

S No.	Chapter Title	Page No.
1	Number System	1
2	LCM & HCF	7
3	Ratio & Proportion	9
4	Percentage	12
5	Mensuration	20
6	Time & Work	29
7	Speed Time & Distance	31
8	Simple Interest	34
9	Compound Interest	37
10	Profit & Loss	43
11	Average	46
12	Algebra	53
13	Geometry	59
14	Trigonometry	80
15	Statistics (Measures of Central Tendency)	89
16	Age Problems	94
17	Pipe & Cistern	99
18	Data Interpretation	102

1 CHAPTER

Number System



Complex Number (Z)

$Z = \text{Real numbers} + \text{Imaginary numbers}$

$$Z = a + ib$$

Where, $a = \text{Real numbers}$.
 $b = \text{Imaginary numbers}$.

Real Numbers

Rational and irrational numbers together are called real numbers. These can be represented on the number line.

Imaginary Numbers

Numbers that can not be represented on the number line.

Integer Numbers

A set of numbers which includes whole numbers as well as negative numbers, is called integer numbers, it is denoted by I.

$$I = \{-4, -3, -2, -1, 0, 1, 2, 3, 4, \dots\}$$

Natural Numbers

The numbers which are used to count things are called natural numbers.

$$N = \{1, 2, 3, 4, 5, \dots\}$$

Whole Numbers

When 0 is also included in the family of natural numbers, then they are called whole numbers.

$$W = \{0, 1, 2, 3, 4, 5, \dots\}$$

The product of four consecutive natural numbers is always exactly divisible by 24.

Even Numbers

Numbers which are completely divisible by 2 are called even numbers.

$$n^{\text{th}} \text{ term} = 2n$$

$$\text{Sum of first } n \text{ even natural numbers} = n(n+1)$$

Sum of square of first n even natural

$$\text{numbers} = \frac{2n(n+1)(2n+1)}{3}$$

$$\left\{ n = \frac{\text{Last term}}{2} \right\}$$

Odd Numbers

The numbers which are not divisible by 2 are odd numbers.

$$\text{Sum of first } n \text{ odd numbers} = n^2$$

$$\left\{ n = \frac{\text{Last term} + 1}{2} \right\}$$

Natural Numbers

$$\text{Sum of first } n \text{ natural numbers} = \frac{n(n+1)}{2}$$

$$\begin{aligned} \text{Sum of square of first } n \text{ natural numbers} \\ = \frac{n(n+1)(2n+1)}{6} \end{aligned}$$

Sum of cube of first n natural numbers =

$$\left[\frac{n(n+1)}{2} \right]^2$$

The difference of the squares of two consecutive natural numbers is equal to their sum.

$$\text{Example - } 11^2 = 121$$

$$12^2 = 144$$

$$11 + 12 \rightarrow 23$$

$$\text{Difference } 144 - 121 = 23$$

Prime Numbers – Which have only two forms - $1 \times$ numbers

$$\text{E.g. - } \{2, 3, 5, 7, 11, 13, 17, 19, \dots\}$$

Where, 1 isn't a Prime Number.

- The digit 2 is only even prime number.
- 3, 5, 7 is the only pair of consecutive odd prime numbers.
- Total prime numbers between 1 to 25 = 9
- Total prime numbers between 25 to 50 = 6
- There are total of 15 prime numbers between 1-50.
- There are total of 10 prime numbers between 51 – 100.
So there are total 25 prime numbers from 1-100.
- Total prime numbers from 1 to 200 = 46
- Total prime numbers from 1 to 300 = 62
- Total prime numbers from 1 to 400 = 78
- Total prime numbers from 1 to 500 = 95

Co-prime Numbers

Numbers whose HCF is only 1.

$$\text{E.g. - } (4,9), (15, 22), (39, 40)$$

$$\text{HCF} = 1$$

Perfect Number

A number whose sum of its factors is equal to that number (except the number itself in the factors)

$$\text{E.g. - } 6 \rightarrow 1, 2, 3 \rightarrow \text{Here } 1 + 2 + 3 \rightarrow 6$$

$$28 \rightarrow 1, 2, 4, 7, 14 \rightarrow 1 + 2 + 4 + 7 + 14 \rightarrow 28$$

Rational Numbers

Numbers that can be written in the form of P/Q , but where Q must not be zero and P and Q must be integers.

$$\text{E.g. - } 2/3, 4/5, \frac{10}{-11}, \frac{7}{8}$$

Irrational Numbers

These cannot be displayed in P/Q form.

$$\text{E.g. - } \sqrt{2}, \sqrt{3}, \sqrt{11}, \sqrt{19}, \sqrt{26} \dots$$

Perfect square numbers



Unit Digit which can be of square

0
1
4
5 or 25
6
9

Which can't be square

2 —
3 —
7 —
8 —

- The last two digits of the square of any number will be the same as the last two digits of the square of numbers 1-24.

Note: Therefore, everyone must remember the squares of 1-25.

Convert to Binary and Decimal –

1. Convert Decimal Number to Binary Number

To find the binary number equivalent to a decimal number, we continuously divide the given decimal number by 2 until we get 1 as the final quotient.

E.g.

2	89	$2 \times 44 = 88 ; 89 - 88 = 1$
2	44	$2 \times 22 = 44 ; 44 - 44 = 0$
2	22	$2 \times 11 = 22 ; 22 - 22 = 0$
2	11	$2 \times 5 = 10 ; 11 - 10 = 1$
2	5	$2 \times 2 = 4 ; 5 - 4 = 1$
2	2	$2 \times 1 = 2 ; 2 - 2 = 0$
	1	Final quotient

Hence, binary number equivalent to 89 = $(1011001)_2$

2. Convert Binary to Decimal Number

In binary system the value of 1 when it moves one place to its left every time it doubles itself and wherever 0 comes its value is 0.

E.g.

1	0	1	1	0	0	1
2^6	2^5	2^4	2^3	2^2	2^1	2^0

Now

$$\begin{aligned} (1011001)_2 &= 1 \times 2^6 + 0 \times 2^5 + 1 \times 2^4 + 1 \times 2^3 + 0 \times 2^2 + 0 \times 2^1 + 1 \times 2^0 \\ &= 64 + 0 + 16 + 8 + 8 + 0 + 1 \quad \{2^0 = 1\} \\ &= 89 \end{aligned}$$

Finding the Number of Divisors or Number of Factors

First we will do the prime factorization of the number and write it as Power and multiply by adding

One to each power, then the number of divisors will be obtained.

Ex: By how many total numbers can 2280 be completely divided?

Sol. $2280 = 2^3 \times 3^1 \times 5^1 \times 19^1$

$$\begin{aligned} \text{Number of divisors} &= (3 + 1) (1 + 1) (1 + 1) (1 + 1) \\ &= 4 \times 2 \times 2 \times 2 = 32 \end{aligned}$$

Find the unit's digit

1. When the number is in the form of power –

When the unit digit of Base is 0, 1, 5 or 6, the unit digit of the result remains the same for any natural power. When the unit digit of base is 2, 3, 4, 7, 8, or 9, divide the power by 4 and put the same power on the unit digit of the base as the remainder. When the power is rounded off to 4, then the 4th power will be placed on the unit digit of the base.

2. In the form of simplification –

Write the unit digit of each number and simplify it according to the symbol, the result that will come will be its unit digit answer.

Divide by Power of Numbers (Finding the Divisor)

- If $a^n + b^n$ is given –
If n is odd, then $(a+b)$ will be its divisor.
- If $a^n - b^n$ is given –
Divisor (when n is odd) $\rightarrow (a-b)$
Divisor (when n is even) $\rightarrow (a - b)$ or $(a + b)$ or both.

- If $a^n \div (a - 1)$ then the remainder always be 1.
- $a^n \div (a + 1)$ $\left\{ \begin{array}{l} \text{If } n \text{ is an even then the remainder always be 1.} \\ \text{If } n \text{ is an odd then the remainder always be } a. \end{array} \right.$
- If $(a^n + a) \div (a - 1)$ then the remainder always be 2 .
- $(a^n + a) \div (a + 1)$ $\left\{ \begin{array}{l} \text{If } n \text{ is an even then the remainder always be zero (0).} \\ \text{If } n \text{ is an odd then the remainder always be } (a - 1) \end{array} \right.$

Terminating Decimal

Those numbers which end after a few digits after the decimal like - 0.25, 0.15, 0.375 can be written in a fraction number.

Non-Terminating Decimal

Those numbers which continue after the decimal and can be of two types.

0.3333, 0.7777, 0.183183183.....

Repeating

Numbers that never end after the decimal, but repeat, till infinity. It can be written in fractions.

Non Repeating Decimal

Numbers that never end after the decimal point, but they do not repeat their numbers.

Recurring Decimal Fraction

That decimal fraction is the repetition of one or more digits after the decimal point, then one or more digits are repeated after the dot.

Eg. $\frac{1}{3} = 0.333\dots$, $\frac{22}{7} = 3.14285714\dots$. To represent such fractions, a line is drawn over the repeating digit.

$$0.\overline{3524} = \frac{3524 - 35}{9900} = \frac{3489}{9900} = \frac{1163}{3300}$$

$$\frac{22}{7} = 3.14285714\dots = 3.\overline{142857}$$

It is called bar.

- Convert pure recurring decimal fraction to simple fraction as follows –

$$0.\overline{p} = \frac{p}{9} \quad 0.\overline{pq} = \frac{pq}{99} \quad 0.\overline{pqr} = \frac{pqr}{999}$$

- Convert a mixed recurring decimal fraction to an ordinary fraction as follows –

$$0.p\overline{q} = \frac{pq - p}{90} \quad 0.pq\overline{r} = \frac{pqr - pq}{900}$$

$$0.p\overline{q}r = \frac{pqr - p}{990} \quad 0.pqrs\overline{r} = \frac{pqrs - pq}{9900}$$

Example -

$$(i) 0.\overline{39} = \frac{39}{99} = \frac{13}{33}$$

$$(ii) 0.\overline{625} = \frac{625 - 6}{990} = \frac{619}{990}$$

$$(iii) 0.\overline{3524} = \frac{3524 - 35}{9900} = \frac{3489}{9900} = \frac{1163}{3300}$$

Symbol of the Roman Method

1	→	I
2	→	II
3	→	III
4	→	IV
5	→	V
6	→	VI
7	→	VII
8	→	VIII
9	→	IX
10	→	X
20	→	XX
30	→	XXX
40	→	XL
50	→	L
100	→	C
500	→	D
1000	→	M

Rule of Divisibility

Rule of 2	The last digit is an even number or zero (0) as - 236, 150, 1000004
Rule of 3	If the sum of the digits of a number is divisible by 3, then the whole number will be divisible by 3. E.g. 729, 12342, 5631
Rule of 4	Last two digits are zero or divisible by 4. E.g. 1024, 58764, 567800
Rule of 5	The last digit is zero or 5. E.g. 3125, 625, 1250
Rule of 6	If a number is divisible by both 2 and 3 then it is also divisible by 6. E.g. 3060, 42462, 10242
Rule of 7	After multiplying the last digit of a number by 2 and

	subtracting it from the remaining number, if the number is a multiple of 0 or 7 or if any digit is repeated in a multiple of 6, then the number will be divisible by 7. E.g. 222222, 444444444444, 7854
Rule of 8	If the last three digits of a number are divisible by 8 or the last three digits are '000' (zero). E.g. 9872, 347000
Rule of 9	If the sum of the digits of a number is divisible by 9, then the whole number will be divisible by 9.
Rule of 10	The last digit should be zero (0).
Rule of 11	If the difference between the sum of digits at odd places and sum of digits at even places is zero (0) or 11 or a multiple of 11. E.g. 1331, 5643, 8172659
Rule of 12	Composite form of divisible by 3 and 4.
Rule of 13	Repeating the digit 6 times, or multiplying the last digit by 4 and adding it to the remaining number, if the number is divisible by 13, then the whole number will be divisible by 13. E.g. 222222, 17784

Practice Questions

- Q.1** If $\frac{3}{4}$ of a number is 7 more than $\frac{1}{6}$ of that number, then what will be $\frac{5}{3}$ of that number?
 (a) 12 (b) 18
 (c) 15 (d) 20
- Q.2** If the sum of two numbers is a and their product is b then their reciprocals will be –
 (a) $\frac{1}{a} + \frac{1}{b}$ (b) $\frac{b}{a}$
 (c) $\frac{a}{b}$ (d) $\frac{a}{ab}$
- Q.3** The sum of two numbers is 75 and their difference is 25, then what will be the product of those two numbers?
 (a) 1350 (b) 1250
 (c) 1000 (d) 125
- Q.4** Divide 150 into two parts such that the sum of their reciprocal is $\frac{3}{112}$. Calculate both parts.
 (a) 50, 90 (b) 70, 80
 (c) 60, 90 (d) 50, 100
- Q.5** If the sum of any three consecutive odd natural numbers is 147, then the middle number will be –
 (a) 47 (b) 48
 (c) 49 (d) 51

- Q.6** If the product of first three and last three of 4 consecutive prime numbers is 385 and 1001, then find the greatest prime number.
- Q.7** What will be the sum of the even numbers between 50 and 100?
- Q.8** What will be the sum of odd numbers between 50 and 100?
- Q.9** In a division method, the divisor is 12 times the quotient and 5 times the remainder. Accordingly, if the remainder is 36, then what will be the dividend?
 (a) 2706 (b) 2796
 (c) 2736 (d) 2826
- Q.10** What is the unit digits of $(3694)^{1739} \times (615)^{317} \times (841)^{491}$
 (a) 0 (b) 2
 (c) 3 (d) 5
- Q.11** What will be written in the form of $\frac{p}{q}$ of 18.484848....?
 (a) $\frac{462}{25}$ (b) $\frac{610}{33}$
 (c) $\frac{200}{11}$ (d) $\frac{609}{33}$
- Q.12** Put $\frac{0.936 - 0.568}{0.45 + 2.67}$ in the form of rational number.
- Q.13** What will be the common factor of $\{(127)^{127} + (97)^{127}\}$ and $\{(127)^{97} + (97)^{97}\}$?
 (a) 127 (b) 97
 (c) 30 (d) 224

Answer Key

- | | | | |
|----------------|----------------|-----------------|---|
| Q.1 (d) | Q.2 (c) | Q.3 (b) | Q.4 (b) |
| Q.5 (c) | Q.6 13 | Q.7 1800 | Q.8 1875 |
| Q.9 (c) | Q.10(a) | Q.11(b) | Q.12 $\frac{2024}{17205}$ |
| Q.13(d) | | | |

2

CHAPTER

Least Common Multiple and Highest Common Factor (LCM & HCF)

Factor: A number is said to be a factor of another if it completely divides the other. Like 3 and 4 are factors of 12.

Common Factor: The number which completely divides two or more given numbers is called the common factor of those numbers. Thus, one common factor of 9, 18, 21 and 33 is 3.

LCM (Least common multiple)

- The smallest number which is completely divisible by the given numbers is called LCM.
- **Finding the LCM of the number having power** - After factoring the prime, we will write it in the form of quotient and the number of primes that will be used will be written as multiplication and will keep the maximum power on it.

Ex-1: Find LCM of $(12)^{16}$, $(18)^{15}$, $(30)^{18}$

Sol. $(12)^{16} = (2 \times 2 \times 3)^{16} = (2^2 \times 3)^{16} = 2^{32} \times 3^{16}$
 $(18)^{15} = (2 \times 3 \times 3)^{15} = (2 \times 3^2)^{15} = 2^{15} \times 3^{30}$
 $(30)^{18} = (2 \times 3 \times 5)^{18} = 2^{18} \times 3^{18} \times 5^{18}$
 Therefore, LCM = $2^{32} \times 3^{30} \times 5^{18}$ Ans.

LCM of fractions

$$\text{LCM} = \frac{\text{LCM of Numerator}}{\text{HCF of Denominator}}$$

Ex-2: Find LCM of $\frac{1}{2}$ and $\frac{5}{8}$?

Sol. - $\text{LCM} = \frac{\text{LCM of 1 and 5}}{\text{HCF of 2 and 8}} \Rightarrow \frac{5}{2}$

HCF (Highest Common Factor)

- The greatest number by which all the given numbers are completely divisible is called HCF.
- Like – H.C.F. of 18 and 24 is 6.

Ex.1: If the H C F of two numbers is found by the division method, then the quotient is 3, 4, and 5 respectively. If the mean of two numbers is 18, then find the numbers.

Sol. - There are two numbers a and b

$$\begin{array}{r} a \overline{) b} \quad 3 \\ \quad c \overline{) a} \quad 4 \\ \quad \quad d \overline{) c} \quad 5 \\ \quad \quad \quad x \quad x \end{array}$$

The last denominator is HCF.

$$d = 18$$

$$c = 5 \times d = 5 \times 18 = 90$$

$$a = (4 \times c) + d$$

$$= (4 \times 90) + 18 = 378$$

$$b = 3a + c$$

$$= (3 \times 378) + 90 = 1134 + 90 = 1224$$

So, the numbers are 1224 and 378

To find the HCF of a number with powers-

- First factor it into the base and write it as a power, and write it as a multiplication of all prime numbers in the base and put the lowest power on it.

Ex:1 Find HCF of $(24)^8$, $(36)^{12}$, $(18)^{16}$

Sol. $24 = (2^3 \times 3)^8 = 2^{24} \times 3^8$
 $36 = (2^2 \times 3^2)^{12} = 2^{24} \times 3^{24}$
 $18 = (2 \times 3^2)^{16} = 2^{16} \times 3^{32}$
So, HCF = $2^{16} \times 3^8$

Finding the HCF of a Fraction –

$$\text{HCF} = \frac{\text{HCF of Numerator}}{\text{LCM of Denominator}}$$

Ex: $\frac{18}{25}, \frac{12}{7}, \frac{6}{35}$

Sol. $\frac{\text{HCF of 18, 12, 16}}{\text{LCM of 25, 7, 35}}$

- HCF of Addition of any two numbers and their L.C.M is equal to the HCF of given two numbers.

Let the two numbers be x and y, and their H.C.F is H.

Therefore, $x = Ha$

$$y = Hb$$

Where a and b are mutually prime.

LCM of x, y = Hab

$$\text{and } x + y = H(a + b)$$

Now 'a' and 'b' are mutually prime numbers, then $(a + b)$ and ab will also be primes with each other. So we can conclude that the HCF of $H(a + b)$ and Hab is H which is also the H.C.F of x and y.

Relation between LCM and HCF: -

$\text{LCM} \times \text{HCF} = \text{Product of both numbers}$

Ex.1 The LCM and HCF of two numbers are 420 and 28. If one number is 84, find the other number –

Sol. Second Number = $\frac{420 \times 28}{84} = 140$

- The smallest number for x, y, z in which the remainder r is left after dividing, The answer for this will be (LCM of x, y, z + r).

Practice Questions

- Q.1** What is the greatest common factor of 84, 126, 140 ?
- Q.2** Find HCF of $x^6 - 1$ and $x^4 + 2x^3 - 2x^1 - 1$
(a) $x^2 + 1$ (b) $x - 1$
(c) $x^2 - 1$ (d) $x + 1$
- Q.3** What will be the LCM of 15, 18, 24, 27, 36?
- Q.4** Six bells started ringing simultaneously, if these bells rang respectively at an interval of 2, 4, 6, 8, 10, 12 seconds, then how many times in 30 minutes will they ring together?
(a) 4 times
(b) 10 times
(c) 16 times
(d) None of the above
- Q.5** Three persons start walking together on a 11 km long circular path in the same direction. Their speed is 4, 5.5 and 8 km/hr respectively. After how much time will they meet together at the starting point ?
- Q.6** Find the greatest common factor of 1.75, 5.6 and 7.
- Q.7** The sum of two numbers is 36, their highest common factor is 3 and the least common factor is 105, what will be the sum of the reciprocals of these numbers ?
(a) $\frac{2}{35}$ (b) $\frac{3}{25}$
(c) $\frac{4}{35}$ (d) $\frac{2}{25}$
- Q.8** Find two such three-digit numbers whose L.C.M. is 5760 and H.C.F. is 80?

Answer Key

- 1. 14** **2. (c)** **3. 1080**
4. (c) **5. 22 Hours** **6. 0.35**
7. $\frac{4}{35}$ **8. 640 and 720**

3

CHAPTER

Ratio & Proportion

Ratio

The ratio of two quantities is that fraction by which another quantity can be expressed in terms of one quantity.

Ratio - Proportion is a comparison of two quantities by part. The ratio of **a** to **b** can be written in the following way.

$$a : b = \frac{a}{b} = a \div b$$

Mixed Ratio

The product of the previous terms of two or more ratios and the new ratio formed by the product of the last terms is called mixed ratio.

Ex. Mixed ratio of 4 : 3, 9 : 13, 26 : 5, 2 : 15

$$\frac{4 \times 9 \times 26 \times 2}{3 \times 13 \times 5 \times 15} = \frac{16}{25}$$

Inverse or Inverse Ratio

The ratio in which the increase of the first quantity decreases the quantity of the second is called inverse proportion.

Inverse ratio of a : b

$$= \left(\frac{1}{a} : \frac{1}{b} \right) \times (\text{LCM of a and b})$$

Combined Ratio

1. If the ratio between the first and second quantities = a : b

And, between the second and third quantities = c : d

Then the combined ratio between these three quantities.

$$\begin{array}{c} a : b \\ \swarrow \quad \searrow \\ \quad c : d \\ \hline ac : bc : bd \end{array}$$

Ex. If A : B = 4 : 5 and B : C = 6 : 7 then A : C = ?

Sol.

$$\begin{array}{c} A : B : C \\ 4 : 5 \\ \swarrow \quad \searrow \\ \quad 6 : 7 \\ \hline 24 : 30 : 35 \end{array}$$

Hence, A : C = 24 : 35

2. Ratio between the first and second quantities = a : b

Ratio between the second and third quantities = c : d

Ratio between the third and fourth quantities = e : f

$$\begin{array}{c} a : b \\ \swarrow \quad \searrow \\ \quad c : d \\ \quad \quad \searrow \\ \quad \quad \quad e : f \\ \hline ace : bce : bde : bdf \end{array}$$

Ex. If A : B = 1 : 2, B : C = 3 : 4, C : D = 2 : 3 then,

A : B : C : D = ?

Sol.

$$\begin{array}{c} A : B : C : D \\ 1 : 2 \\ \quad 3 : 4 \\ \quad \quad 2 : 3 \\ \hline \end{array}$$

$$\underline{\underline{6 : 12 : 16 : 24 \text{ or } 3 : 6 : 8 : 12}}$$

Proportion

In both $6 : 18$ and $8 : 24$, 6 is one third of 18 and 8 is one third of 24. This equality of ratios is called proportion.

a, b, c, d are called terms. **a** is the first term, **b** is the second, **c** is the third and **d** is the fourth term.

- a, d are external terms.
- b, c middle terms.
- d is the fourth proportional.
- From the expression $a : b : c : d$
 1. First proportional = $a = b^2/c$
 2. Mid proportional = $b = \sqrt{ac}$
 3. Third proportional = $c = \frac{b^2}{a}$

Ex. What will be the first proportional of 6 and 9?

Sol. $a = \frac{b^2}{c} = \frac{6^2}{9} = \frac{36}{9} = 4$

Ex. What will be the proportion between 0.32 and 0.02?

Sol. $b = \sqrt{ac} \Rightarrow \sqrt{0.32 \times 0.02} = \sqrt{0.0064} \Rightarrow 0.08$ Ans.

Characteristics of Ratio

1. Alternendo

$$\text{If } \frac{a}{b} = \frac{c}{d} \text{ then } \frac{a}{c} = \frac{b}{d}$$

2. Componendo

If $a : b :: c : d$

Then, $(a + b) : b :: (c + d) : d$

$$\text{i.e., } \frac{a}{b} = \frac{c}{d}$$

$$\text{Thus, } \frac{a+b}{b} = \frac{c+d}{d}$$

3. Dividendo

If $a : b :: c : d$ then

$$(a - b) : b :: (c - d) : d$$

$$\text{i.e., } \frac{a}{b} = \frac{c}{d}$$

$$\text{Thus, } \frac{a-b}{b} = \frac{c-d}{d}$$

4. Compondo & Dividendo

It includes sum-ratio and inter-ratio.

If $a : b :: c : d$ is in proportion.

Then, $(a + b) : (a - b) :: (c + d) : (c - d)$

$$\text{or, } \frac{a+b}{a-b} = \frac{c+d}{c-d}$$

If x is divided between P and Q in the ratio $a : b$, then

$$P\text{'s part} = \frac{a}{a+b} \times x$$

$$Q\text{'s part} = \frac{b}{a+b} \times x$$

Difference of parts of P and

$$Q = \frac{a-b}{a+b} \times x \quad (\text{Where, } a > b)$$

If x liter mixture of milk and water contains milk and water in the ratio $a : b$, then to make the ratio $c : d$ of milk and water in that mixture, $\frac{x(ad-bc)}{c(a+b)}$ liters of water will have to be added.

Practice Question

Q.1 The ratio of the length and breadth of a field is $5:2$. If the width of the field is 40 metres, then find the length of the field?

- (a) 200 meters (b) 100 meters
(c) 50 meters (d) 80 meters

Q.2 Two natural numbers are in the ratio $3 : 5$ and their product is 2160. Which of these will be the smaller number?

- (a) 36 (b) 24
(c) 8 (d) 12

Q.3 Three numbers are in the ratio 3 : 4 : 5 respectively. If the sum of the first and the third number is 52 more than the second number, then what is the greatest number?

- (a) 65 (b) 52
(c) 79 (d) 63
(e) None of these

Q.4 The ratio of difference, sum and product of two numbers is 1 : 7 : 24, what will be the product of these numbers?

- (a) 36 (b) 52
(c) 48 (d) 40

Q.5 What is the third proportional of 16 and 24?

- (a) 20 (b) 18
(c) 36 (d) 40

Q.6 The ratio of last year's income of A, B, C is 3 : 4 : 5, their individual previous year and this year's income are in the ratio 4 : 5, 2 : 3 and 3 : 4 respectively. If the sum of the present income of all three is Rs.78,800, then find the income of each of them this year?

Q.7 A bag contains coins of Rs. 1, 50 paise and 25 paise in the ratio 5 : 6 : 8. If there is a total of 210 rupees in this bag, then find the number of coins of each type?

Q.8 The present ratio of the ages of two boys is 5 : 6. After 2 years this ratio becomes 7 : 8. Then what will be the ratio after 12 years?

- (a) $\frac{22}{24}$ (b) $\frac{15}{16}$
(c) $\frac{17}{18}$ (d) $\frac{11}{12}$

Q.9 A man divided his property in such a way that the shares of his daughter and wife and that of his wife and son are each in the ratio of 3 : 1. If the son got Rs. 10,000 less than the daughter, then what is the total value of the property?

- (a) Rs. 15,250 (b) Rs. 16,250
(c) Rs. 17,500 (d) Rs. 18,500

Q.10 Rs. 600 is distributed among A, B and C in such a way that Rs 40 more than $\frac{2}{5}$ of A, Rs 20 more than $\frac{2}{7}$ of B and Rs 10 more than $\frac{9}{17}$ of C are mutually equal, in this, what is the share of A?

- (a) 150 (b) 170
(c) 200 (d) 280

Answer Key

- | | | | |
|--|----------------|-----------------|----------------|
| Q.1 (b) | Q.2 (a) | Q.3 (a) | Q.4 (c) |
| Q.5 (c) | | | |
| Q.6 A = 18000, B = 28800 & C = 32000 | | | |
| Q.7 Number of 1 rupee coins = $(5 \times 21) = 105$
Number of 50 paise coins = $(6 \times 21) = 126$
Number of 25 paise coins = $(8 \times 21) = 168$ | | | |
| Q.8 (c) | Q.9 (b) | Q.10 (a) | |

4 CHAPTER

Percentage

- Percentage means 'Per hundred'.
- The fraction whose denominator is 100, is called percentage and the numerator of that fraction is called percentage rate.
- 5 out of 100 = $\frac{5}{100} = 5\%$

- 10 out of 100 = $\frac{10}{100} = 10\%$
- That is, when an amount is compared with 100, it is called percentage. The basis with which the comparison is made. In the fraction, the base will be called the denominator.

Conversion From Percentage to Fraction

$100\% = 1$	$10\% = \frac{1}{10}$	$5\frac{5}{19}\% = \frac{1}{19}$
$50\% = \frac{1}{2}$	$9\frac{1}{11}\% = \frac{1}{11}$	$5\% = \frac{1}{20}$
$33\frac{1}{3}\% = \frac{1}{3}$	$8\frac{1}{3}\% = \frac{1}{12}$	$4\frac{1}{6}\% = \frac{1}{24}$
$25\% = \frac{1}{4}$	$7\frac{9}{13}\% = \frac{1}{13}$	$4\% = \frac{1}{25}$
$20\% = \frac{1}{5}$	$7\frac{1}{7}\% = \frac{1}{14}$	$2\frac{1}{2}\% = \frac{1}{40}$
$16\frac{2}{3}\% = \frac{1}{6}$	$6\frac{2}{3}\% = \frac{1}{15}$	$37\frac{1}{2}\% = \frac{3}{8}$
$14\frac{2}{7}\% = \frac{1}{7}$	$6\frac{1}{4}\% = \frac{1}{16}$	$62\frac{1}{2}\% = \frac{5}{8}$
$12\frac{1}{2}\% = \frac{1}{8}$	$5\frac{15}{17}\% = \frac{1}{17}$	$57\frac{1}{7}\% = \frac{4}{7}$
$11\frac{1}{9}\% = \frac{1}{9}$	$5\frac{5}{9}\% = \frac{1}{18}$	$66\frac{2}{3}\% = \frac{2}{3}$
$80\% = \frac{4}{5}$	$75\% = \frac{3}{4}$	

Note -

- To convert a fraction or a decimal fraction or an integer into a percentage, multiply it by 100.
- To convert a percentage to a fraction, divide it by 100.

- One number is what percent of another number –

$$\% \text{ Quantity} = \frac{\text{Given number}}{\text{Basic (another number)}} \times 100$$

Ex.1 What percent of 48 is 6 ?

Sol. Let x % of 48 = 6

$$\text{Now, } 48 \times \frac{x}{100} = 6$$

$$\frac{12x}{25} = 6$$

$$x = \frac{6 \times 25}{12} = 12 \frac{1}{2}$$

The percent $12 \frac{1}{2}$ % of 48 is 6.

Ex.2 Convert $\frac{9}{16}$ into percentage.

Sol. $\frac{9}{16} = \left(\frac{9}{16} \times 100 \right) \% = \frac{225}{4} \% = 56 \frac{1}{4} \%$

Ex.3 When 60 is subtracted from 60% of a number, the result is 60. The number is-

- (a) 120 (b) 150
(c) 180 (d) 200

Sol. (b)

Let the number be x, then –

$$X \times 60\% - 60 = 60$$

$$\Rightarrow 0.60x - 60 = 60$$

$$\Rightarrow 0.60x = 120$$

$$\Rightarrow x = \frac{120}{0.60}$$

$$X = 200$$

∴ The number is 200.

If there is a percentage change

Case I – If there is increase of x_1 % & x_2 % then –

$$\text{Percentage increase} = x_1 + x_2 + \frac{x_1 \cdot x_2}{100}$$

Case II – If there is decrease of x_1 % & x_2 % then –

$$\text{Percent decrease} = x_1 + x_2 - \frac{x_1 \cdot x_2}{100}$$

Case III – If there is increase of x_1 % and decrease of x_2 % then –

$$\text{Percentage change} = x_1 - x_2 - \frac{x_1 \cdot x_2}{100} \left[\begin{array}{l} x_1 = \text{Always percentage increase} \\ x_2 = \text{Always percentage decrease.} \end{array} \right]$$

Note- If you want to calculate the percentage change in expenses, sales income, revenue, area, etc., then the same rule will be used.

Ex.1 Two successive price increases of 10% and 10% of an article are equal to which one single price increase?

- (a) 19% (b) 20%
(c) 21% (d) 22%

Sol. (c)

Let the initial price of any item = 100

New price = 110 % of 100 of 110%.

$$\left(100 \times \frac{110}{100} \times \frac{110}{100} \right) = 121 = 21\%$$

Hence, the single price increase is equal to 21 percent.

Ex.2 The price of laptop is increased by 25%. Now by what percent is the price increased for the second time so that the total increase becomes 35%?

- (a) 7.5 (b) 9
(c) 8 (d) 10

Sol. (c)


Given that –

Price increased by 25% = x
 The overall increase was 35%.
 Formula:
 Total Increment = $x + y + \frac{xy}{100}$
 Calculation:
 $\Rightarrow 35 = 25 + y + (25 \times y/100)$
 $\Rightarrow 35 = 25 + y + y/4$

$\Rightarrow 140 = 100 + 4y + y$
 $\Rightarrow 140 - 100 = 5y$
 $\Rightarrow 40 = 5y$
 $\Rightarrow y = 8$
 Hence, the second time increase is 8%.

Method II

Let the price of laptop = 100



$$\text{Percent increase} = \frac{135 - 125}{125} \times 100$$

$$= \frac{10}{125} \times 100 = 8\%$$

Ex.3 The value of a tool decreases by 20% every year. What will be the cost of that tool after 3 years?

- (a) 48.8% (b) 51.2% (c) 54% (d) 60%

Sol. (a)

20% = 1/5

Initial	Final
5	4
5	4
5	4
125	64

Required Percentage = $\frac{64}{125} \times 100 = 48.8\%$

Method II

Let the price of tool = x

According to question,

$$\text{Price after 8 years} = x \times \frac{80}{100} \times \frac{80}{100} \times \frac{80}{100}$$

$$= \frac{64}{125} x$$

$$\text{Decrease in price} = x - \frac{64}{125}x = \frac{61}{125}x$$

$$\text{Percentage decrease} = \frac{\frac{61}{125}x \times 100}{x} = 48.8\%$$

Formula Related to Population

- If the population of a city is P and it is increasing at the rate of x% per annum, then the population after n years –

$$= P \left(1 + \frac{x}{100} \right)^n$$

- If it is decreasing, then the population

$$= P \left(1 - \frac{x}{100} \right)^n$$

- Ex.1** If the population of a city is increasing at the rate of 4% per annum and the present number of people is 15625, then what will be the population after 3 years?

Sol. Required Population = $15625 \left(1 + \frac{4}{100} \right)^3$
 $= 15625 \times \frac{26}{25} \times \frac{26}{25} \times \frac{26}{25} = 17576$

- Ex.2** The population of a city is 8000. If the number of males increases at the rate of 6 percent and the rate of increase in the number of females is 10 percent, then the population will become 8600. Find the number of females in the city.

Sol. Let the number of females = x
 Then, 110 % of x + 106% of (8000 + x)
 = 8600
 $\frac{110x}{100} + \frac{106(8000 + x)}{100} = 8600$
 $x(110 + 106) = 8600 \times 100 - 8000 \times 106$

$$\therefore x = \frac{8600 \times 100 - 8000 \times 106}{110 + 106}$$

$$= \frac{12000}{4} = 3000$$

- Ex.3** In 1988 the population of a city decreased by 12%. In 1989 it increased by 15%. What was the overall impact on the city's population in the early 1990s?

Sol. % Impact = % Increase – % Decrease –
 $\frac{(\% \text{ Increase})(\% \text{ Decrease})}{100}$

$$= 15 - 12 - \frac{15 \times 12}{100} = 15 - 13.8 = 1.2$$

\therefore The population increased by 1.2%.

- If x percent of any amount is taken by the first person, y percent of the balance is taken by the second person and after taking z percent of the remaining amount by the third person, if A is left, then the total amount initially –

$$= \frac{A \times 100 \times 100 \times 100}{(100 - x)(100 - y)(100 - z)}$$

- An increase or decrease in the price of a commodity, a decrease or increase in its consumption –

(a) % Increase in consumption
 $= \frac{100 \times \text{decrease}}{100 - \text{decrease}}$

(b) % Decrease in consumption
 $= \frac{100 \times \text{increase}}{100 + \text{increase}}$