



**RRRB - JE**



**CIVIL**

**Railway Recruitment Board**

Volume - 1

**Engineering Mechanics**



# CONTENTS

S.No.	Topic	Page No.
1.	Introduction .....	1 – 1
2.	System of Forces .....	2 – 9
3.	Friction & Application .....	10 – 17
4.	Center of Gravity .....	18 – 25
5.	Moment of Inertia .....	26 – 32
6.	Kinematics .....	33 – 43
7.	Trusses .....	44 – 54
8.	Collisions .....	55 – 62
9.	Rigid Body Dynamics in Plane Motion .....	63 – 70
10.	Objective Sheet .....	71 – 87

# SYSTEM OF FORCES

## THEORY

### 2.1 | DEFINITIONS

**Force** - It is something that have either displacement or deformation effect on the body

**Line of action of force** - It is the direction of effect on the body. It is also the direction of force

### 2.2 | LAW OF TRANSMISSIBILITY OF FORCE

Along the line of action if force is shifted, effect of force will remain same




### 2.3 | SYSTEM OF FORCES


When multiple forces act on the body in different direction, at different point and of different magnitude, it is called system of forces

It can be of two types:

- Coplanar Forces (2D Forces)
- Non Coplanar forces (3D Forces)




**Concurrent Forces :** If the line of action of all the force meet at a single point, then it is called concurrent for system.




$T_1, T_2$  &  $W$  are Meeting at 'O'

**Non concurrent forces :** If the line of action of all forces do not meet at a single point, then it is called non-concurrent force system.



**Parallel Forces :** if the line of action of all forces do not meet anywhere, then it is called parallel force system.




**Collinear Forces :** If all the forces are having same line of action, then is is called collinear force system.




## 2.4 | FREE BODY DIAGRAM

It is the diagram representing all the external forces acting on a body or the system of bodies.


### Example 1




F.B.D. For  $W_1$





F.B.D. For  $W_2$



F.B.D. For  $W_3$




### Example 2



**2.5 | LAW OF FORCES**


**2.5.1 Law of Parallelogram**



$$F = \sqrt{F_1^2 + F_2^2 + 2F_1F_2 \cos\theta}$$

$$\tan \alpha = \frac{F_2 \sin\theta}{F_1 + F_2 \cos\theta}$$

**Case-I**




$\Rightarrow$  Collinear force

$$F = F_1 + F_2$$

**Case-II**




$\Rightarrow$  Perpendicular force




$$F = \sqrt{F_1^2 + F_2^2}$$

$$\tan \alpha = \frac{F_2}{F_1}$$

**Case-III**




$\Rightarrow$  Collinear force



$$F = F_1 - F_2 \quad [F_1 > F_2]$$

$$F = F_2 - F_1 \quad [F_2 > F_1]$$


### 2.5.2 Lami's Theorem (Sine Rule)



For three concurrent forces to be in equilibrium

$$\Rightarrow \frac{F_1}{\sin \theta_1} = \frac{F_2}{\sin \theta_2} = \frac{F_3}{\sin \theta_3}$$

### 2.5.3 Component of Forces



In this method forces are resolved along two perpendicular axes.

$$F_x = F_1 \cos \theta_1 - F_2 \cos \theta_2 - F_3 \sin \theta_3 + F_4 \cos \theta_4$$

$$F_y = F_1 \sin \theta_1 + F_2 \sin \theta_2 - F_3 \sin \theta_3 - F_4 \sin \theta_4$$


For resultant force

$$F = \sqrt{(\sum F_x)^2 + (\sum F_y)^2}$$


$$\tan \theta = \frac{F_y}{F_x}$$

### Example 3

Determine reaction by inclined plane on the sphere



**Solution:**




$$\frac{T}{\sin 120^\circ} = \frac{R}{\sin 120^\circ} = \frac{1000}{\sin 120^\circ}$$

$$T = R = 1000 \text{ N}$$


**Example 4**

A string of length 10m supporting a weight of 120 N as shown in fig.



Determine Tension in string

**Solution:**



$$2T \cos \theta = 120$$


$$T = \frac{120 \times 5}{2 \times 4}$$

$$T = 75 \text{ N}$$



**Example 5**

With what acceleration 'a' should the box of figure descend so that the block of mass  $m$  exerts a force  $Mg/4$  on the floor of the box?

**Solution:**

The block is at rest with respect to the box which is accelerated with respect to the ground. Hence, the acceleration of the block with respect to the ground is 'a' downward. The force on the block are

- (i)  $Mg$  downward (by the earth) and
- (ii)  $N$  upward (by the floor).

The equation of motion of the block is, therefore,

$$Mg - N = Ma.$$

If  $N = Mg/4$ , the above equation gives  $a = 3g/4$ . The block and hence the box should descend with an acceleration  $3g/4$ .

**Example 6**

Two bodies of masses  $m_1$  and  $m_2$  are connected by a light string going over a smooth light pulley at the end of an incline. The mass  $m_1$  lies on the incline and  $m_2$  hangs vertically. The system is at rest. Find the angle of the incline and the force exerted by the incline on the body of mass  $m_1$ .



**Solution:**

Figure shows the situation with the forces on  $m_1$  and  $m_2$  shown. Take the body of mass  $m_2$  as the system. The forces acting on it are



- (i)  $m_2g$  vertically downward (by the earth),
- (ii)  $T$  vertically upward (by the string).

As the system is at rest, these forces should add to zero.

This gives  $T = m_2g$ . ... (i)

Next, consider the body of mass  $m_1$  as the system. The forces acting on this system are

- (i)  $m_1g$  vertically downward (by the earth),
- (ii)  $T$  along the string up the incline (by the string)
- (iii)  $N$  normal to the incline (by the incline).

As the string and the pulley are all light and smooth, the tension in the string is uniform everywhere. Hence, same  $T$  is used for the equations of  $m_1$  and  $m_2$ . As the system is in equilibrium, these forces should add to zero.

Taking components parallel to the incline,

$$T = m_1g \cos\left(\frac{\pi}{2} - \theta\right) = m_1g \sin\theta. \quad \dots(ii)$$

Taking components along the normal to the incline,

$$N = m_1g \cos\theta \quad \dots(iii)$$

Eliminating  $T$  from (i) and (ii),

$$m_2g = m_1g \sin\theta$$

or,  
giving

$$\sin\theta = m_2/m_1$$

$$\theta = \sin^{-1}(m_2/m_1).$$

From (iii)

$$N = m_1g \sqrt{1 - (m_2/m_1)^2}$$

□□□

Unleash the topper in you